

Unbiased Geothermal Temperature Gradient = GradTGN

[@wikipedia](#)

Rate of [Unbiased Geothermal Temperature Profile](#) $T_{GN}(\mathbf{r})$ change along [True Vertical Direction](#) z :

$$(1) \quad \text{Grad}T_{GN}(\mathbf{r}) = \frac{\partial T_{GN}}{\partial z}$$

It can be estimated as ratio of [true vertical component](#) of regional [Earth's Heat Flux](#) $j_z(\mathbf{r})$ to the [subsurface Thermal Conductivity](#) $\lambda_e(\mathbf{r})$ at a certain [subsurface point](#) $\mathbf{r}$:

$$(2) \quad \text{Grad}T_{GN}(\mathbf{r}) = \frac{j_z(\mathbf{r})}{\lambda_e(\mathbf{r})}$$

The most common value of [Unbiased Geothermal Temperature Gradient \(GT\)](#) away from tectonic plate boundaries is:
 $G_T = 0.025 \div 0.035 \text{ } ^\circ\text{C/m}$.

The difference between [Geothermal Temperature Gradient \(GradTG\)](#) and [Unbiased Geothermal Temperature Gradient \(GradTGN\)](#) is that the former accounts for the effects of the seasonal/daily temperature variations, which are only significant above [Neutral Temperature Layer](#).

See Also

[Geology / Geothermal Temperature Field](#)

[[Geothermal Temperature Gradient \(GradTG\)](#)] [[Geothermal Temperature Profile \(TG\)](#)]